

## MATH CIRCLE: SEQUENCES AND SERIES

LAMC OLYMPIAD GROUP

Let us look at the numbers  $1, \frac{1}{2}, \frac{1}{3}, \dots$ . This sequence seems to be approaching zero, and yet at no point is it actually equal to zero. So how do we formalize this notion?

**Definition.** A *sequence* of real numbers is an assignment  $a : \mathbf{N} \rightarrow \mathbf{R}$ , i.e. an ordered set of real numbers  $a_1, a_2, a_3, \dots$  (we may also write  $\{a_n\}_{n \geq 1}$  or  $\{a_n\}$  for this sequence). Such a sequence is said to *converge* to a real number  $L$ , called the *limit* of the sequence, if and only if for all offsets  $\varepsilon > 0$ , there is an integer  $N$  depending on  $\varepsilon$  such that

$$L - \varepsilon < a_n < L + \varepsilon, \text{ for all } n > N.$$

We write  $\lim_{n \rightarrow \infty} a_n = L$ .

**Problem 1.** Show that if a sequence  $\{a_n\}$  converges to both  $L$  and  $K$ , then in fact  $L = K$ . That is, the limit is unique if it exists.

**Problem 2.** Show that a convergent sequence is bounded. That is, if  $\{a_n\}$  converges to some real number  $L$ , then there exists a constant  $A$  such that  $-A < a_n < A$  for all  $n \geq 1$ .

**Problem 3. (a)** Show that if  $\{a_n\}$  and  $\{b_n\}$  are convergent sequences with  $a_n \leq b_n$ , then

$$\lim_{n \rightarrow \infty} a_n \leq \lim_{n \rightarrow \infty} b_n.$$

(Note that if  $a_n < b_n$ , we can still *only* conclude that  $\lim_{n \rightarrow \infty} a_n \leq \lim_{n \rightarrow \infty} b_n$ .)

**(b)** Show that if  $\{a_n\}, \{b_n\}, \{c_n\}$  are sequences with  $a_n \leq b_n \leq c_n$ , such that  $\{a_n\}$  and  $\{c_n\}$  converge to the same limit, then  $\{b_n\}$  also converges to that limit.

**Problem 4.** Show that if  $\{a_n\}$  converges to  $L$  and  $\{b_n\}$  converges to  $K$ , then  $\{a_n + b_n\}$  converges to  $L + K$  and  $\{a_n \cdot b_n\}$  converges to  $L \cdot K$ . (It is also true that if  $a_n > 0$  converge to  $L$  then  $a_n^r$  converge to  $L^r$  for  $r \in \mathbf{R}$ .)

**Definition.** The *series* associated to a sequence  $\{a_n\}$  is the new sequence  $\{s_n\}$  of *partial sums* given by

$$s_n = a_1 + \dots + a_n.$$

We say that the series converges to a real number  $S$  if  $s$  converges  $S$ . In that case, we write  $\sum_{n=1}^{\infty} a_n = S$  and call this limit the *value* of the series. If the series doesn't converge, we say that it *diverges*.

**Problem 5.** Show that if the series  $\sum_n a_n$  converges, then the sequence  $\{a_n\}$  must converge to 0. (*Hint: subtract two consecutive partial sums and use Problem 4.*)

**Problem 6. (a)** Show that if  $-1 < r < 1$ , then  $\sum_{n=0}^{\infty} r^n = \frac{1}{1-r}$ .

**(b)** We toss a fair coin until we hit heads on the  $n$ th try. What is the probability that  $n$  is even?

**(c)** Interpreting the base-10 number  $0.a_1a_2a_3 \dots$  as  $\sum_{n \geq 1} \frac{a_n}{10^n}$ , show that  $0.999 \dots = 1$ .

**Axiom.** Any increasing bounded sequence is convergent.

(To show this we would need to formally construct the real numbers; this is one of their defining features. Note that this doesn't work if we only allow rational numbers.)

**Problem 7.** Show the same statement for decreasing sequences.

**Problem 8.** Let  $\{a_n\}$  be the sequence defined recursively by  $a_1 = 1$  and  $a_{n+1} = \sqrt{2 + a_n}$ . Show that  $\lim_{n \rightarrow \infty} a_n = 2$ .

**Problem 9.** Given two sequences  $\{a_n\}$  and  $\{b_n\}$  of non-negative integers such that  $a_n \leq b_n$  and  $\sum_n b_n$  converges, show that  $\sum_n a_n$  converges.

**Definition.** A subsequence of a sequence  $\{a_n\}$  is given by selecting only some terms of the sequence. That is, it is given by  $\{a_{k_n}\}$  where  $\{k_n\}$  is a strictly increasing sequence of indices  $k_1 < k_2 < \dots$ .

**Problem 10.** Show that any subsequence of a convergent sequence converges to the same limit.

**Problem 11.** Given any two sequences  $\{a_n\}$  and  $\{b_n\}$  both converging to the same number  $L$ , show that the new sequence

$$c_n = \begin{cases} a_{(n+1)/2} & \text{if } n \text{ is odd} \\ b_{n/2} & \text{if } n \text{ is even} \end{cases}$$

given by interlacing  $a$  and  $b$  also converges to  $L$ .

**Problem 12. (a)** Show that  $\sum_n \frac{1}{n}$  diverges.

**(b)** Show that  $\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = 1$ .

**(c)** Show that  $\sum_n \frac{(-1)^n}{n}$  and  $\sum_n \frac{1}{n^2}$  converge.

**Problem \*13.** Suppose that  $\sum_n a_n$  and  $\sum_n b_n$  are two series with the same terms but in a different order, and assume that all  $a_n \geq 0$ . Show that if  $\sum_n a_n$  converges then  $\sum_n b_n$  converges to the same limit.

**Problem \*14.** Show that any bounded sequence has a convergent subsequence.

**Problem \*15.** Let  $S$  be a nonempty bounded set of real numbers. Show that there is real number  $u$  such that  $x \leq u$  for all  $x \in S$ , and which is minimal with this property. This is called a supremum, denoted  $u = \sup S$ .