

# Hyperbolic Geometry II - Angles in the Hyperbolic Plane

Yan Tao

ORMC Advanced 1 - Winter 2023

## 1 Hyperbolic Triangles


This week, we continue the example of the *hyperbolic plane* that we started with last week. Last week, we showed that in the hyperbolic plane, given any line and a point not on that line, there are infinitely many parallel lines through that point. We now continue developing this new geometry by studying shapes - the most basic of which is the *triangle* (a shape with three hyperbolic line segments for sides). (Technically, the most basic shape would have two sides, but that is impossible in the hyperbolic plane - think about why.) As a reminder:

**Definition 1** *In Poincaré's model of the hyperbolic plane, also known as the **Poincaré disk**:*

- *The points are all points on the plane that are less than 1 away from the origin in distance.*
- *The points on the unit circle  $S^1$  (exactly 1 away from the origin in distance) are called the **ideal points**.*
- *The (hyperbolic) lines are given by circles orthogonal to  $S^1$ , as well as diameters of  $S^1$ . The angles between shapes in general is the angle between their tangent lines at the intersection point.*

**Problem 1** *Show that the angles of a (non-degenerate) triangle add up to less than 180 degrees. (Hint: Last week, we learned that circle inversions preserve circles and angles. Use an inversion to move some sides of the triangle to diameters, as it is easier to find the angle between straight lines.)*

As it turns out, the angles of a hyperbolic triangle depend on the side lengths of the triangle. Those familiar with Euclidean geometry will recall a similar relationship in the plane. Before we quantify this relationship, let's look at it qualitatively. In the following diagrams, points  $A$  and  $C$  are the same points, while points  $B, B',$  and  $B''$  lie on the same ray from point  $A$ . Each diagram shows a triangle, so  $AC, AB, AB',$  and  $AB''$  are diameters.



**Problem 2** Like in the Euclidean case, we define "Angle  $B$ " to be the angle at vertex  $B$  (so in this case, angle  $ABC$ ). Which of the angles  $B, B',$  and  $B''$  is largest? Smallest? Is there a pattern?

Recall the hyperbolic distance function defined previously:

**Definition 2** Let  $A$  and  $B$  be points in the hyperbolic plane. Let  $P$  and  $Q$  be the ideal endpoints of the hyperbolic line connecting  $A$  and  $B$  such that  $P$  is closer to  $B$  and  $Q$  is closer to  $A$ . The **hyperbolic distance** from  $A$  to  $B$  is given by

$$d(A, B) := \log([A, B; P, Q])$$

where the cross ratio is given by

$$[A, B; P, Q] = \frac{AP \cdot BQ}{AQ \cdot BP}$$

using the Euclidean distances in that formula.

**Problem 3** Fix  $A$  inside the disk, and take a point  $B$  approaching the boundary. Show that  $d(A, B)$  gets larger and larger as you do this (we say that  $d(A, B) \rightarrow \infty$ ).


This suggests a natural extension of the distance function we've defined.

**Definition 3** If  $A$  or  $B$  is an ideal point, we say that  $d(A, B) = \infty$ .

**Problem 4** Let  $ABC$  be a hyperbolic triangle with  $d(A, B) = d(B, C) = \infty$ . What is angle  $B$ ?

## 2 The Euclidean Functions sin and cos

In the Euclidean case, there is likewise a relationship between the sides of a triangle and its angles. To quantify this, we use *trigonometric functions*, which are first defined for acute angles.



**Definition 4** The functions *sine* and *cosine* of the angle  $\theta$  are defined by

$$\sin(\theta) = \frac{\text{opposite}}{\text{hypotenuse}} = \frac{a}{c} \text{ and } \cos(\theta) = \frac{\text{adjacent}}{\text{hypotenuse}} = \frac{b}{c}$$


**Problem 5** Calculate:

- $\cos(30^\circ)$
  
  
- $\sin(45^\circ)$
  
  
- $\cos(60^\circ)$

**Problem 6** Let  $\theta$  be an acute angle. Show that

$$\cos^2(\theta) + \sin^2(\theta) = 1$$

Problem 6 shows that the functions sin and cos lie on the unit circle, like so:



This leads us to define the sine and cosine of any angle:

**Definition 5** The *cosine* of any angle is the x-coordinate of the intersection of the ray from the origin at that angle with the unit circle. The *sine* is the y-coordinate of that intersection.

The third important trig function, which we will not study this week, is the *tangent*, which is just sine divided by cosine. The cosine function in particular allows us to find side lengths from angles, or vice versa.

**Problem 7** Calculate:

- $\cos(135^\circ)$
  
  
- $\sin(180^\circ)$
  
  
- $\cos(240^\circ)$

**Theorem 1** (*Euclidean Law of Cosines*) Let  $A, B, C$  be the angles at the respective vertices and  $a, b, c$  be the (Euclidean) side lengths of the side lengths opposite to the respective vertices. Then


$$\cos(A) = \frac{b^2 + c^2 - a^2}{2bc}$$

**Problem 8** (*Bonus*) Prove Theorem 1.

**Problem 9** Consider increasing  $a, b, c$ . We know by the triangle inequality that we better increase two at a time, so consider all the ways to do that. What happens to angle  $A$  as you keep increasing  $a$  and  $b$ ? What about  $a$  and  $c$ ? And finally,  $b$  and  $c$ ? (*Hint: Consider all the possible angles of a triangle - is cosine increasing on this domain, decreasing, or neither?*)

### 3 The Functions sinh and cosh

Just like the functions sine and cosine are defined by rays intersecting the circle  $x^2 + y^2 = 1$ , we can define the *hyperbolic sine and cosine* (abbreviated sinh and cosh respectively) by rays intersecting the hyperbola  $x^2 - y^2 = 1$ . This is illustrated below, but is unwieldy to calculate this way, so we'll use a formula (which is equivalent to finding the rays) instead.



**Definition 6** Given any real number  $a$ , the functions  $\sinh(a)$  and  $\cosh(a)$  are given by

$$\cosh(a) = \frac{e^a + e^{-a}}{2} \text{ and } \sinh(a) = \frac{e^a - e^{-a}}{2}$$

Additionally, just like with the Euclidean sine and cosine, the **hyperbolic tangent** is defined by  $\tanh(a) = \sinh(a)/\cosh(a)$ .

**Problem 10** When is  $\cosh(a) > 0$ ? When is  $\sinh(a) > 0$ ?

**Problem 11** Find the range of  $\tanh$ .

**Problem 12** Show that  $\tanh$  is a bijection, and find  $\tanh^{-1}$ . (Hint: Recall that any function with an inverse that's defined for all values in the range is a bijection. To find the inverse, set  $\tanh(y) = x$  and solve for  $y$ .)

**Problem 13** Show that in the hyperbolic plane, given any point  $A$  which is Euclidean distance  $r$  from the origin  $O$ , the hyperbolic distance satisfies

$$d(O, A) = 2 \tanh^{-1}(r)$$

## 4 Angles of Hyperbolic Triangles

Just like in the Euclidean case, we use the convention that angles  $A, B, C$  of hyperbolic triangle  $ABC$  are at the corresponding vertices, and  $a, b, c$  are the hyperbolic side lengths opposite vertices  $A, B, C$ , respectively. We'll make use of the following theorem:

**Theorem 2** (*Hyperbolic Law of Cosines*)

$$\cos(A) = \frac{\cosh(b) \cosh(c) - \cosh(a)}{\sinh(b) \sinh(c)}$$

**Problem 14** *Prove Theorem 2.*

**Problem 15** *Like in problem 9, consider increasing  $b$  and  $c$ . (This corresponds to moving vertex  $A$  towards the boundary.) What happens to angle  $A$ ? Compare to your answer to Problem 4.*



**Problem 16** (*Challenge*) *What happens when you increase  $a$  and  $b$  as in Problem 9?*

**Problem 17** *In Problem 1, we showed that the sum of angles of any hyperbolic triangle are less than 180 degrees. How does the sum of angles depend on the side lengths?*