

## NUMBER BASES

MATH CIRCLE (ADVANCED) 10/21/2012

We write the number three thousand six hundred fifty seven as

$$3657 = 3 \cdot 1000 + 6 \cdot 100 + 5 \cdot 10 + 7 = 3 \cdot 10^3 + 6 \cdot 10^2 + 5 \cdot 10^1 + 7 \cdot 10^0.$$

Is there anything special about using 10? From a mathematical point of view, the answer is no.

**Proposition:** Fix some  $n$  (called the base). We can write any number uniquely in the form

$$a_k n^k + a_{k-1} n^{k-1} + \cdots + a_2 n^2 + a_1 n + a_0 n^0$$

where each  $a_i$  takes values from 0 to  $n - 1$ .

To avoid ambiguity, we will write the base as a subscript. For example, in base 6, the number fifty ( $50_{10}$ ) is  $1 \cdot 6^2 + 2 \cdot 6 + 2 \cdot 6^0 = 122_6$ .

1) Convert the following numbers to base 10.

a)  $1232_4$

b)  $10120_3$

c)  $723_9$

d)  $100111_2$

e)  $1232_5$

---

2) Convert the number  $98_{10}$  into: (can you come up with a fast way to do this?)

a) base 7

b) base 4

c) base 3

d) base 2

3) Write out multiplication tables for base 2 and base 3.

We can add/multiply numbers using the same methods we know for base 10.

4) Calculate (in the same base as given):

a)  $11010_2 + 10011_2$

b)  $21012_3 + 121201_3$

c)  $1001_2 \cdot 101_2$

d)  $2111_3 \cdot 122_3$

5) Is it possible that the following statements are true in some number base system?  
Is so, what base?

a)  $3 \cdot 4 = 10$

b) both  $3 + 4 = 10$  and  $3 \cdot 4 = 15$

c) both  $2 + 3 = 5$  and  $2 \cdot 3 = 11$

6) State and prove a condition (involving the representation of a number) which allows us to determine whether the number is even or odd:

a) in the base 3 system.

b) in the base  $n$  system.

7) An evil king wrote three secret two-digit numbers  $a, b, c$ . A handsome prince must name three numbers  $X, Y, Z$ , after which the king will tell him the sum  $aX + bY + cZ$ . The prince must then name all three of the King's numbers, or he will be executed. Help out the prince!

8)\* Prove the proposition stated at the beginning of the handout.

9)\* Prove that from the set  $0, 1, 2, \dots, 3^k - 1$  one can choose  $2^k$  numbers so that none of them can be represented as the arithmetic mean of some pair of the chosen numbers.

Some problems are taken from:

- D. Fomin, S. Genkin, I. Itenberg “Mathematical Circles (Russian Experience)”